

# LANADELUMAB

BRAND NAME	TAKHZYRO
DRUG CLASS	Monoclonal antibody (human) for hereditary angioedema (HAE)
AVAILABILITY	Prefilled syringe contains 300 mg/2 mL of lanadelumab. Also contains dibasic sodium phosphate dihydrate, citric acid monohydrate, histidine, sodium chloride and polysorbate-80. <sup>1</sup> Contain 3.45 mg (0.15 mmol) of sodium. <sup>1</sup> The solution is clear to slightly opalescent and colourless to slightly yellow. <sup>1</sup>
pH	6.0 <sup>1</sup>
PREPARATION	Ready to use. Allow the syringe to reach room temperature. <sup>2</sup> <b>Do not shake.</b> <sup>1</sup>
STABILITY	Store at 2 to 8 °C. Do not freeze. Protect from light. <sup>1</sup> Stable for 14 days below 25 °C. Do not return to the fridge. <sup>1</sup>
ADMINISTRATION	
<b>IM injection</b>	Not recommended
<b>SUBCUT injection</b>	Inject into the abdomen, thigh or upper arm. Rotate the site of injection. <sup>1</sup> The first dose must be given under medical supervision. Subsequent doses may be suitable for self-administration after appropriate training. <sup>1</sup>
<b>IV injection</b>	Not recommended
<b>IV infusion</b>	Not recommended
COMPATIBILITY	No information
INCOMPATIBILITY	No information
SPECIAL NOTES	Injection site reactions are usually mild to moderate and include pain, flushing, redness, itching and rash. <sup>1</sup>

## REFERENCES

1. Product information. Available from [www.tga.gov.au](http://www.tga.gov.au). Accessed 24/03/2022.
2. Consumer medicine information. Available from [www.tga.gov.au](http://www.tga.gov.au). Accessed 24/03/2022.