

CARBETOCIN

BRAND NAME	DURATOCIN, CARBETOCIN WOCKHARDT
DRUG CLASS	Oxytocin analogue
AVAILABILITY	Duratocin ampoule contains 100 microgram/mL of carbetocin. Also contains methionine, succinic acid, mannitol and sodium hydroxide. ¹ Carbetocin Wockhardt prefilled syringe contains 100 microgram/mL of carbetocin. Also contains sodium chloride and glacial acetic acid. ¹ The solution is clear and colourless. ^{1,2}
pH	Duratocin: 5.5 ¹ Carbetocin Wockhardt: 3.9 ³
PREPARATION	Not required
STABILITY	Duratocin: store below 30 °C ¹ Carbetocin Wockhardt: store at 2 to 8 °C. Do not freeze. Protect from light. ¹
ADMINISTRATION	
IM injection	Suitable for Duratocin, after a vaginal delivery. ¹
SUBCUT injection	Not recommended
IV injection	Suitable after a caesarean section or after a vaginal delivery. ¹ Inject slowly over 1 minute. ¹
IV infusion	Not recommended
COMPATIBILITY	No information
INCOMPATIBILITY	No information
SPECIAL NOTES	Carbetocin is only given after the delivery of the infant. ¹

REFERENCES

1. Product information. Available from www.tga.gov.au. Accessed 15/06/2022.
2. Consumer medicine information. Available from www.tga.gov.au. Accessed 15/06/2022.
3. Gray D. Carbetocin Wockhardt prefilled syringes pH [email]. Manly, NSW: Interpharma Pty Ltd; 15/06/2022.